

Characterization of epiphytic ferns *Asplenium indicum* By GC-MS From Panchgani and Mahabaleshwar area of Maharashtra

Damayanti Choudhari, Manda Ghatage

Department of Botany Kisan Veer Mahavidyalaya Wai

Department of Botany, B V's M.B. S. K., Kanya Mahavidyalaya, Kadegaoan, Sangali, India.

Abstract : The GCMS analysis of methanolic extracts of *Asplenium indicum* leaves revealed the presence of some compounds. The compound includes Ethanol, 1-(2-butoxyethoxy)-, 3-Allyl-6-methoxyphenol, Ethanol, 2-(2-butoxyethoxy)-acetate, Neophytadiene, Hexanedioic acid, bis (2-ethylhexyl) ester, Hexadecanoic acid, 2-hydroxyl-1-(hydroxymethyl) ethyl ester. Ethanol, 2-(2-butoxyethoxy)-acetate. From the results concluded that the epiphytic fern *Asplenium indicum* leaves contain various bioactive compounds which have various medicinal properties.

Keywords: *Asplenium indicum*, epiphytic, 3-Allyl-6-methoxyphenol, bioactive.

Introduction:

Pteridophytes are vascular cryptogams epiphytic, shade loving from the dominant vegetation on the earth (Fraser Jenkins 2008). These plant groups have an important role in ecosystem and had various ethnomedicinal uses which had to be investigated in detail. Unfortunately pteridophytes are ignored groups of plants and had significant medicinal properties which are traditionally used by ethnic communities (Annie and Kumaresan 2010, Vashishta *et al.*, 2012). *Asplenium indicum* is an epiphytic pteridophyte found in hilly region of Western Ghat of India. In recent years Gas chromatography – Mass spectroscopy has been applied as a technique to identify the structures of different bioactive components from plant extracts with great success (Prasain 2004, De Rijke 2006). Hence in the present investigation was carried out to determine the possible bioactive compounds of *Asplenium indicum* by GC-MS studies.

Methods:

GC-MS analysis was conducted to identify the chemical composition of the one epiphytic fern species under study. Analysis was carried out on a GC Clarus 500 Perkin Elmer system comprising an AOC-20i auto sampler and gas chromatograph interfaced to a mass spectrometer. Column Elite -1 fused silica capillary column operating in electron impact mode at 70 eV. For analysis Helium was used as carrier gas. [Continuous flow of 1ml/min.]. 2µl extract prepared in methanol was injected to the instrument Turbo gold mass detector and

turbomass 5.1 software detected the sample during 36 min of the extraction process. Oven was kept at 110⁰C with 2 min holding temperature of injector was kept at 250⁰C. Other important parameters include- 1. Mass spectra were taken at 17EV, 2. A scan interval of 0.5 S 3. Fragmented from 45 to 450 Da the MS detection was completed in 38 min. By comparing average peak area of component with the total area, relative percentage of amount of each component was calculated. Biological activities of the known components of the spectrum were determined by using literature.

Result:

GC-MS analysis of *Aspleniumindicum* accommodate the occurrence of various bioactive phytoconstituents in the methanolic extracts. The bioactive principles with their RT, molecular formula, molecular weight and concentration (%), biological activity in the methanolic extracts of *Aspleniumindicum* was presented in the (Fig. and Table1).

Fig 1. GC-MS analysis of *Aspleniumindicum*




Table 1. GCMS peak report of *Aspleniumindicum*.

Name of the compound	R.T	Peak Area	Molecular Formula	Molecular Weight	Biological Activity
Ethanol,1-(2-butoxyethoxy)-	9.04 2	1236 7	C ₈ H ₁₈ O ₃	162	-

3-Allyl-6-methoxyphenol	13.1 22	6113 3	$C_{10}H_{12}O_2$	164	Antihistaminic, anti-inflammatory and Antioxidant activity (Hussein 2016) antiseptic and anesthetic, analgesic, antioxidant, anti-inflammatory and antimicrobial activities(Jadhav 2004)
Ethanol,2-(2-butoxyethoxy)-acetate	13.2 88	7614 4	$C_{10}H_{20}O_4$	204	paints, adhesives, plastics, and paint remover(Yang 2020)
Neophytadiene	19.7 75	3755 7	$C_{20}H_{38}$	278	Antipyretic , analgesic acid, anti-inflammatory, antioxidant, antimicrobial.(Adnan 2019), (Raman 2012)
Hexanedioic acid, bis(2-ethylhexyl)ester	28.8 38	2907 1	$C_{22}H_{42}O_4$	370	Antioxidant, antimicrobial, ant proliferative (Nguyen2018). Acidifier, acidulant, antiuric acid production (Arora 2018) (Dr.Duke's Phytochemical and Ethno botanical Database)
Hexanedioic acid,2-hydroxy-1hydroxymethyl)ethyl ester	30.4 97	1183 5	$C_{19}H_{38}O_4$	330	Hemolytic, pesticide, flavor, antioxidant (Arora and Kumar 2018)

Discussion and conclusion:

Karthishwaran *et al.*, (2012) Concluded that GCMS are important technique for separation of compounds. These compounds showed pharmacological properties. Bertzet *et al.*, (1997) reported that GCMS has many applications including it is the best technique for identification and separation of volatile and semi-volatile compounds. Previous studies on GC-MS analysis showed the occurrence of various compounds in *Adiantum tetraphyllum* (Melos *et al.*, 2007), *Salvinia molesta* (Choudhary *et al.*, 2008), *Marsilea quadrifolia* (Sivagurunathan and Xavier, 2014) and *Actinopteris radiata* (Manonmani and Sara, 2015). The present study showed various bioactive compounds with pharmacological properties.

Janakiraman (2015) observed major bioactive components 2methylbutance,4-diol, 3-(l-ethoxyethoxy)-, 2-hydroxy-5-methylbenzaldehyde, 3-5 dinydroxy-4-methylbenzaldehyde. *Dryanariaquercifolia* revealed the presence of major compounds namely Tetrahydroisovelleral 7, 10, Pentadeadi-Yanoic acid, Phosphoric acid, Octyldiphenyl Ester, Octicizer Phosphoric acid, 2ethylhexyl Diphenyl Ester, Quercetin 7, 3, 4 trimethox, 30triacotanediol Ergost-s-en3-0l. Compounds are reported s by (Kalpna Devi R 2016).

The results of GC-MS analysis showed the presence of different phytoconstituents in the studied *Aspleniumindicum* species with varied retention time. The identified phytochemicals from the fern under study showed anti-inflammatory, anti-uric acid, anti-oxidant, anti-bacterial, hepatoprotectant, antifungal, Antihistaminic, antiseptic and anesthetic, analgesic, antipyrepic, analgesic acid, hemolytic, pesticide, , Further phytochemical and pharmacological analysis of the epiphytic fern *Aspleniumindicum* under study may use as natural drugs developments.

References:

Adnan, M.D., Chy, N.U. and Kamal, M. (2019). Investigation of the Biological Activities and Characterization of Bioactive Constituents of Ophiorrhizarugosa var. prostrata (D.Don) and Mondal Leaves through In Vivo, In Vitro, and In Silico Approaches: *Molecules*,**24**:13-67.

Annie, R. and Kumaresan, V. (2010).Pteridophytes, Gymnosperms and Palaeobotany, 3rd edn, Saras Publication, Nagercoil,:1-4.

Arora, S. and Kumar, G. (2018).Screening of bioactive compounds from leaf of *cenchrusciliarisl*.From that region of Rajasthan, India.*Ijpsr*, **9**(5):1878-1885.

Bertz, J.M., Gay, M.L., Mossoba, M.M., Adams, S. and Portz, B.S. (1997).Chiral gas chromatographic determination of ephedrine-type alkaloids in dietary supplements containing Ma Huang.*Journal of AOAC International*,**80**:303-315.

Choudhary, M.I., Naheed, N., Abbaskhan, A., Musharraf, S.G., Siddiqui, H. and Rahman, A.U. (2008).Phenolic and other constituents of fresh water fern *Salviniamolesta*.*Phytochemistry*; **69**(4):1018-1023.

De Rijke, E., Out, P., Neissen, W.M.A., Ariese, F., Gooijer, C., and Brinkman, U.A. (2006).The analytical separation and detection methods for flavonoids. *Journal of Chromatography*, 1112, 31–63.

Fraser Jenkins, C.R. (2008). Taxonomic revision of three hundred Indian subcontinentalpteridophytes with a revised census-list-a new picture of fern taxonomy and nomenclature in the Indian subcontinent.Bishen Singh Mahendra Pal Singh, Dehradoun, India.

Hussein, H.M., Hameed, H.I. and Ibraheem, O.A. (2016). Antimicrobial Activity and Spectral Chemical Analysis of Methanolic Leaves Extract of *AdiantumCapillus-Veneris* Using GC-MS and FT-IR Spectroscopy, *International Journal of Pharmacognosy and Phytochemical Research*, **8**(3):369-385.

Jadhav, B.K., Khandelwa, I K.R., Ketkar, A.R. and Pisal, S.S. (2004). Formulation and evaluation of mucoadhesive tablets containing eugenol for the treatment of periodontal diseases. *Drug Development and Industrial Pharmacy*.**30**(2):195-203.

Janakiraman, N., Marimuthu., Alias, M. and Johnson A. (2015). GC-MS analysis of ethanolic extracts of *Cyatheanilgirensis* ,*C. gigantean* *C. crinite*. *Egyptain Pharmaceutical Journal*,**15**:43-47.

Kalpana, Devi R., Subramani, V., Annamalai, P., Nakulan, V. and Rajesh, N. J. (2016). Phytochemical analysis, in vitro antioxidant potential and gas chromatography- mass spectrometry studies of *Dicranopterislinearis*. *Asian Journal of Pharmaceutical and Clinical Research*,**9**:1-6.

Karthishwaran, K., Muthukkumarasamy, S. and Sankaran, M. (2012). GCMS analysis of methanolic extract of aerial parts of *Pergulariadaemia*.*Journal of Life Science*,**1**(1):50-55.

Manonmani, R. and Sara, S.C. (2015). GC-MS Analysis of Bioactive Components of an Important Medicinal Fern *Actiniopterisradiata*(Swartz) Link. *World Journal of Pharmaceutical Research*,**4**(4): 1860-1869.

Melos, JLR., Silva, LB., Peres, MTLP., Mapeli, AM., Faccenda, O., Anjos, HH., Torres, TG., Tiviroli, SC., Batista, AL., Almeida, FGN., Flauzino, NS., Tibana, LA., Hess, SC. and Honda NK. (2007). Chemical constituents and evaluation of allelopathic potential of *Adiantumtetraphyllum*Humb.&Bonpl. Ex. Wild (Pteridaceae). *New Chemistry*,**30**:292-297.

Nguyen, T.B., Phan, V. H. L. and Cao, N.D. (2018). Bioactive Compounds from Marine Fungus *Penicilliumcitrinum*Strain ND7c by Gas Chromatography-Mass Spectrometry;*The Pharmaceutical and Chemical Journal*,**5**(1):211-224.

Prasain, J.K., Wang, C.-C., and S. Barnes, S. (2004). Mass spectroscopic methods for the Determination of flavonoids in biological samples. *Free Radical Biology & Medicine*, **37**, 1324–50.

Raman, V. and Samuel, L. (2012). Antibacterial, antioxidant activity and GC-MS analysis of *eupatorium odoratum*: *Asian Journal of pharmaceutical and clinical research*,**5**(2): 99-106.

Sivagurunathan, A. and Xavier, B.I. (2014). GC-MS Evaluation of Bioactive Compounds of *Marsileaquadrifolia*Linn. aquatic Fern. *International Journal for Pharmaceutical Research Scholars*,**3**(1):2277-7873. Sivagurunathan, A. and Xavier, B.I. (2014). GC-MS Evaluation of Bioactive Compounds of *Marsileaquadrifolia*Linn. aquatic Fern. *International Journal for Pharmaceutical Research Scholars*,**3**(1):2277-7873.

Vashishta, P.C.; Sinha, A.K. and Kumar, A. (2012). 29th Edn, Chand and Company Ltd, New Delhi, 1-12.

Yang, J., Chen, J., Huitao, B.I., Haiping, G.U., Zhenling, LIU and Wanxi, P. (2020). Molecules and functions of rosewood, *Pterocarpusindicus*: Molecules and Functions of Rosewood: *PterocarpusIndicusThermal Science*,**24**(3):1869-1876.